

CUTOFF FOR RANDOM TO RANDOM CARD SHUFFLE

MEGAN BERNSTEIN AND EVITA NESTORIDI

ABSTRACT. In this paper, we use the eigenvalues of the random to random card shuffle to prove a sharp upper bound for the total variation mixing time. Combined with the lower bound due to Subag, we prove that this walk exhibits cutoff at $\frac{3}{4}n \log n$, answering a conjecture of Diaconis.

1. INTRODUCTION

One step of the random to random card shuffle consists of picking a card and a position of the deck uniformly, independently at random and moving that card to that position. It was introduced by Diaconis and Saloff-Coste [1] who proved that the mixing time is $O(n \log n)$. Uyemura-Reyes [9] proved that the mixing time is between $\frac{1}{2}n \log n$ and $4n \log n$. Saloff-Coste and Zúñiga [7] improved the upper bound constant by showing that the mixing time is bounded by above by $2n \log n$. Subag [8] proved a lower bound for the mixing time of the form $(\frac{3}{4} - o(1))n \log n$. The best known bound for the total variation distance mixing time is $1.5324n \log n$ is due to Morris and Qin [5]. In this paper we prove that the random to random exhibits cutoff at $\frac{3}{4}n \log n$, which answers a conjecture of Diaconis.

Recently, Dieker and Saliola [3] used representation theory arguments to find the eigenvalues and eigenvectors of this card shuffle. We use this information to give better bounds for the total variation distance, defined as

$$\|P_{id}^{*t} - \pi\|_{T.V.} = \frac{1}{2} \sum_{g \in G} |P_{id}^{*t}(g) - \pi(g)|$$

where $P_{id}^{*t}(g)$ is the probability of starting at the identity and being at g after t steps and π is the uniform measure on S_n . Our first result is:

Theorem 1. *For the random to random card shuffling, if $t = \frac{3}{4}n \log n + cn$ with $c \geq 2$ and n sufficiently large, then*

$$\|P_{id}^{*t} - \pi\|_{T.V.} \leq e^{-c}.$$

Theorem 1 in combination with Subag's lower bound [8] prove the existence of cutoff for the random to random card shuffle. Theorem 1 is proven by bounding the eigenvalues of the random to random card shuffle.

We also consider the following generalization. Let $\nu = [\nu_1, \dots, \nu_m]$ be a partition of n . Then we can perform the random to random to a deck which has ν_i cards of type i , where $i = 1, \dots, m$. We then say that the deck has evaluation ν .

Theorem 2. *For the random to random on a deck with evaluation $\nu = [\nu_1, \dots, \nu_m]$, we have that*

(1) *For $t = \frac{3}{4}n \log n + cn$, then*

$$\|P_{id}^{*t} - \pi\|_{TV} \leq e^{-c}.$$

where $c \geq 2$.

The first author is supported by NSF Grant 1344199.

(2) For $t = \frac{n}{4} \log m + \frac{n}{8} \log n - cn$, then

$$\left\| \frac{P_{id}^{*t}}{\pi} - 1 \right\|_2 \geq \frac{1}{2} e^c.$$

where $c > 0$ and the l^2 norm is with respect to the stationary distribution.

This theorem says that order $n \log n$ steps are necessary and sufficient for the l^2 distance to be small. This is not true for the total variation distance in the case where m is constant. For example, if $\nu = [n-1, 1]$ then an easy coupling argument gives that the mixing time is n , while Theorem 2 says that l^2 mixing time is at least $\frac{n}{8} \log n$.

Although this upper bound falls into the tradition of the pioneering work of Diaconis-Shahshahani, it differs significantly in some respects. Both results use the representation theory of the symmetric group to find a tight upper bound. Since the transposition walk is a conjugacy class walk, in which all elements of a probability class have the same probability of being a generator, the walk can be understood through only the characters of the irreducible representations, with each irreducible representation contributing a single eigenvalue. The random-to-random shuffle is not a conjugacy class walk. The walk, as a result, does not act as a constant on each irreducible representation. This is reflected in the the eigenvalues being indexed by a pair of partitions (λ, μ) where λ is indexing the irreducible representation and μ a further level of decomposition. This leads to the mixing time of the random to random shuffle not being determined solely by the largest eigenvalue and its multiplicity.

This paper is organized as following: Section 2 explains how the eigenvalues of the transition matrix P can be used to bound the l^2 distance. In sections 2.1 and 2.2 we quickly review the work of Diecker and Saliola [3] and we provide bounds for the eigenvalues and their multiplicities. In section 3 we present the proof of Theorem 1. In section 4, we briefly examine the case where the n cards of the deck are not necessarily distinct and prove Theorem 2.

2. THE EIGENVALUES AND THEIR MULTIPLICITIES

The random to random walk is a symmetric, transitive, reversible Markov chain, so its transition matrix diagonalizes with an orthonormal eigenbasis. We now recall Lemma 12.16 of [4] which will allow us to use a l^2 bound.

Lemma 3. *Let P be a reversible, transitive transition matrix of a random walk on a finite state space Ω with eigenvalues $1 = \lambda_1 > \lambda_2 \geq \dots \geq \lambda_{|\Omega|} \geq -1$, then:*

$$4 \|P^t(x, \cdot) - \pi\|_{T.V.}^2 \leq \left\| \frac{P^t(x, \cdot)}{\pi(\cdot)} - 1 \right\|_2^2 = \sum_{j=2}^{|\Omega|} \lambda_j^{2t}$$

for every starting point $x \in \Omega$.

2.1. The eigenvalues. In this section, we will present the formula for the eigenvalues that Diecker and Saliola [3] introduced. We will present some basic tools from the representation theory of the symmetric group, and we will prove a few bounds for the eigenvalues and their multiplicities that will be used to prove Theorem 1.

A partition $\lambda = [\lambda_1, \dots, \lambda_r]$ of n satisfies $\lambda_1 \geq \dots \geq \lambda_r > 0$ and $\lambda_1 + \dots + \lambda_r = n$. We denote $|\lambda| = n$. The Young diagram of λ consists of r left justified rows of cells with λ_i cells in row i . For a partition μ of m , we say μ is contained in λ if $\mu_i \leq \lambda_i$ for each i , and the pair of partitions is called a skew shape. The skew Young diagram λ/μ consists of the cells of λ not in μ . Let λ' and μ' denote the transpose partitions of λ and μ respectively.

The skew Young diagram λ/μ is a horizontal strip if for each i , $\lambda'_i - 1 \leq \mu'_i$, meaning the skew diagram has at most one box per column.

Example 4. If $n = 5$, for $\lambda = \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}$ and $\mu = \begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}$ we have that $\lambda/\mu = \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}$ is a skew diagram and a horizontal strip.

The eigenvalues of the random to random chain are indexed by such pairs (λ, μ) where λ/μ is a horizontal strip. For any Young diagram, let (i, j) denote the cell that is in the i^{th} row and in the j^{th} column. Define the diagonal index of λ :

$$(1) \quad \text{diag}(\lambda) = \sum_{(i,j) \in \lambda} (j - i)$$

Example 5. For $\lambda = \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}$ we have that $\text{diag}(\lambda) = 0 + 1 + 2 - 1 + 0 = 2$.

Diecker and Saliola [3] found that the eigenvalue that corresponds to the pair (λ, μ) is:

$$(2) \quad \text{eig}(\lambda/\mu) = \frac{1}{n^2} \left(\binom{n+1}{2} - \binom{|\mu|+1}{2} + \text{diag}(\lambda) - \text{diag}(\mu) \right)$$

To discuss what the multiplicity of each eigenvalue is, we need to talk about *desarrangement* tableau. Firstly, a *tableau* is a filling of a Young diagram so that each cell is assigned a value. A *standard tableau* of shape λ with $|\lambda| = n$ is one in which the values are from $[n]$ and the entries of each row and column are in strictly increasing order. This necessitates that each entry is used once. An entry i of a tableau is an *ascent* if $i = n$ or if $i < n$ and $i + 1$ occurs weakly to the north and east of i . A standard tableaux is a *desarrangement* tableau if its smallest ascent is even. This is equivalent to either there being no $(1, 2)$ cell of λ or that the value of the $(1, 2)$ cell is odd.

Example 6. The tableau $\mu = \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & \end{array}$ is a *desarrangement* tableaux while $\nu = \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \end{array}$ is not.

Theorem 7 (Dieker-Saliola). *The eigenvalue of the random to random walk corresponding to (λ, μ) is $\text{eig}(\lambda/\mu)$ and occurs with multiplicity $d_\lambda d^\mu$ where d_λ is the number of standard Young tableaux of λ and d^μ is the number of *desarrangement* tableaux of μ .*

Since there are too many eigenvalues to handle individually, it is important in this style of proof to understand how the eigenvalues decay in terms of the partitions. Despite the added complication of two indexing parameters, there is still a partial monotonicity relation on the the eigenvalues for fixed λ as μ changes that is simple to describe. How λ effects the eigenvalue will be established later. The following lemma says that all the eigenvalues of random to random are non-negative.

Lemma 8. *Given Young diagrams λ, μ , with the skew diagram of the difference of λ and μ is a horizontal strip then*

$$(3) \quad \text{eig}(\lambda/\mu) \geq 0$$

Proof. When $\lambda = \mu$, the eigenvalue is zero. For $\lambda \neq \mu$, this will be proven by induction on $n = |\lambda|$.

For $n = 1$, there is only the pair $\lambda = [1]$ and μ the empty partition with $\text{eig}(\lambda/\mu) = 1$. Assume that for n it is true that for every λ, μ , with the skew diagram of the difference of λ and μ is a horizontal strip we have that

$$\text{eig}(\lambda/\mu) \geq 0$$

Then we have that for $n + 1$, assume that λ is a Young diagram with $n + 1$ cells and μ is such that the skew diagram of the difference of λ and μ pair is a horizontal strip. Since $\mu \neq \lambda$ we know that there is there is at least a cell in λ that is not contained in μ . Consider, $\tilde{\lambda}$ to a young diagram that can occur by removing a cell (i, j) from λ that is not included in μ . The induction assumptions says that

$$\text{eig}(\tilde{\lambda}/\mu) \geq 0$$

Now notice then that

$$\text{eig}(\lambda/\mu) = \text{eig}(\tilde{\lambda}/\mu) + \binom{n+2}{2} - \binom{n+1}{2} + j - i = \text{eig}(\tilde{\lambda}/\mu) + n + 1 + j - i \geq 0$$

and this finishes the proof. $\square$

Proposition 9. *Given Young diagrams $\lambda, \mu, \tilde{\mu}$ with the skew diagram of the difference of each λ, μ pair a horizontal strip and $\tilde{\mu} \subseteq \mu$ with $\subseteq$ referring to containment:*

$$\begin{aligned} \text{eig}(\lambda/\mu) &= \text{eig}(\lambda/\tilde{\mu}) - \text{eig}(\mu/\tilde{\mu}) \\ \text{eig}(\lambda/\mu) &\leq \text{eig}(\lambda/\tilde{\mu}) \end{aligned}$$

Proof. Under the given assumptions it is easy to see that $\lambda/\tilde{\mu} = \lambda/\mu \oplus \mu/\tilde{\mu}$ where $\oplus$ means that we just concatenate the cells of the two skew diagrams, getting a new skew diagram. If $\lambda/\tilde{\mu}$ has at most one cell per column then this has to be true for λ/μ and $\mu/\tilde{\mu}$. Therefore, $\mu/\tilde{\mu}$ is a horizontal strip. The second equation occurs from the first, using the fact that $\text{eig}(\mu/\tilde{\mu}) \geq 0$. $\square$

Proposition 9 says that for a fixed λ , its largest eigenvalue corresponds to the smallest μ for which λ/μ is a horizontal strip and there is at least one desarrangement tableau of μ . Let μ_λ be this unique partition. The following proposition gives the eigenvalues corresponding to each (λ, μ_λ) :

Proposition 10. *The largest eigenvalue corresponding to a $(\lambda, \cdot)$ pair, is for (λ, μ_λ) , depends only on λ_1 and λ'_1 , and is:*

$$\text{eig}(\lambda/\mu_\lambda) = \begin{cases} 1 - \frac{(n+1)(n-\lambda_1)}{n^2} & , \text{ if } \lambda'_1 > 2 \\ 1 - \frac{(n+2)(n-\lambda_1)}{n^2} & , \text{ if } \lambda'_1 = 2 \\ 1 & , \text{ if } \lambda'_1 = 1 \end{cases}$$

Proof. If $\lambda'_1 > 2$, μ_λ is given by $\mu'_i = \lambda'_i - 1$. Note that in this case μ_λ is a partition of $n - \lambda_1$. Then, $\text{diag}(\lambda) - \text{diag}(\mu) = \sum_{j=1}^{\lambda_1} j - |\lambda'_j| = \binom{\lambda_1+1}{2} - n$ since λ/μ is a horizontal strip and therefore $(i, j) = (\lambda'_j, j)$. For this case $\text{eig}(\lambda/\mu_\lambda)$ is given by:

$$\begin{aligned} \text{eig}(\lambda/\mu_\lambda) &= \frac{1}{n^2} \left(\binom{n+1}{2} - \binom{n-\lambda_1+1}{2} + \binom{\lambda_1+1}{2} - n \right) \\ &= 1 - \frac{(n+1)(n-\lambda_1)}{n^2} \end{aligned}$$

For the two part partitions $[\lambda_1, n - \lambda_1]$, we have that $\mu_\lambda = [n - \lambda_1, 1]$ where the the cell in the second row of μ_λ is necessary for μ to have a desarrangement tableau. The negative contribution of this extra block is the size of the partition plus the content of the new square, namely $(n - \lambda_1 + 1) + (-1) = n - \lambda_1$.

There are no desarrangement tableaux of a one part partition, so for $\lambda = [n]$ we have that $\mu_{[n]}$ is the empty partition. The content of the partition is $\text{diag}([n]) = 0 + 1 + \dots + (n-1) = \binom{n}{2}$. Then, $\text{eig}([n]) = \frac{1}{n^2} \left(\binom{n+1}{2} + \binom{n}{2} \right) = 1$. $\square$

The following proposition studies the rest of the eigenvalues and provides a bound that will be sufficient in order to get the $\frac{3}{4}n \log n$ bound for the mixing time:

Proposition 11. *For $k > n - \lambda_1$, and μ a partition with λ/μ a horizontal strip and $|\mu| = n - \lambda_1 + k$,*

$$\text{eig}(\lambda/\mu) \leq 1 - \frac{n - \lambda_1}{n} - \frac{k^2 + k(n - \lambda_1)}{n^2}$$

Equality occurs at a partition μ^k that contains μ_λ and the leftmost k boxes of λ/μ_λ .

Proof. The largest eigenvalue for a μ of size $n - \lambda_1 + k$ corresponds to μ given by $\mu'_j = \lambda'_j$ for $j \leq k$ and $\mu'_j = \lambda'_j - 1$ for $j > k$, since the leftmost boxes of λ/μ_λ have the smallest (including possibly negative) content. For the remainder of the proof, let μ refer to this partition. If $k > n - \lambda_1$, $\lambda'_k = 1$. This will make it possible to exactly compute the smallest $\text{eig}(\lambda/\mu)$ over all such μ .

$$\text{eig}(\lambda/\mu) = \frac{1}{n^2} \left(\binom{n+1}{2} - \binom{n - \lambda_1 + k + 1}{2} + \sum_{(i,j) \in \lambda/\mu} j - i \right)$$

Where,

$$\sum_{(i,j) \in \lambda/\mu} j - i = \sum_{j=k+1}^{\lambda_1} j - \lambda'_j = \sum_{j=k+1}^{\lambda_1} j - 1 = \binom{\lambda_1}{2} - \binom{k}{2}.$$

Upon simplification this gives,

$$\text{eig}(\lambda/\mu) = \frac{1}{n^2} (n^2 - (n - \lambda_1)n - (n - \lambda_1)k - k^2),$$

which finishes the proof. $\square$

2.2. Bounding multiplicities. In this section, we introduce bounds on the multiplicities of the eigenvalues. Theorem 7 says that the eigenvalue $\text{eig}(\lambda/\mu)$ occurs with multiplicity $d_\lambda d^\mu$ where d_λ is the number of standard Young tableaux of λ and d^μ is the number of desarrangement tableaux of μ . There is a bijection due to Reiner, Saliola, Welker [6] between tableau of λ and the desarrangement tableau of all partitions μ for which λ/μ is a horizontal strip. As consequence of this, $\sum_\mu d^\mu = d_\lambda$. The total multiplicity of eigenvalues corresponding to a $(\lambda, \cdot)$ pair is thus d_λ^2 .

We will make use of the following proposition which is an outcome of Corollary 2 of Diaconis and Shahshahani [2]. This first bound together with Proposition 10 lead to a short proof of an $n \log n$ upper bound for the mixing time, which we omit.

Proposition 12 (Diaconis-Shahshahani).

$$\sum_{\lambda: n - \lambda_1 = l} d_\lambda^2 \leq \binom{n}{l} \frac{n!}{(n-l)!}$$

The rest of this section is dedicated to proving the following result which will be used to prove the $\frac{3}{4}n \log n$ bound.

We will need one further concept from the combinatorics of tableaux. The *jeu de taquin* is a transformation of a skew standard Young tableau that gives a skew standard Young tableau of a different shape. Given a tableau of a skew shape λ/μ and an empty cell c that can be added to the skew shape, do moves as follows. There are two cells of λ/μ adjacent to c . Move the entry into c that maintains the property that entries are increasing along rows and columns. While the new empty cell is inside the skew shape, continue doing the unique slide that maintains the property. If the empty cell starts on the interior of the skew shape, it moves to the exterior, and vice versa.

Example 13. Starting with the tableau

		2	5
1	3	4	

, *jeu de taquin* on the cell (1, 2) gives:

$$\begin{array}{|c|c|c|c|} \hline & & 2 & 5 \\ \hline 1 & 3 & 4 & \\ \hline \end{array} \rightarrow \begin{array}{|c|c|c|c|} \hline & 2 & & 5 \\ \hline 1 & 3 & 4 & \\ \hline \end{array} \rightarrow \begin{array}{|c|c|c|c|} \hline & 2 & 4 & 5 \\ \hline 1 & 3 & & \\ \hline \end{array} \rightarrow \begin{array}{|c|c|c|c|} \hline & 2 & 4 & 5 \\ \hline 1 & 3 & & \\ \hline \end{array}$$

Example 14. Starting with the same tableau, *jeu de taquin* on the cell (2, 4) gives:

$$\begin{array}{|c|c|c|c|} \hline & 2 & 5 & \\ \hline 1 & 3 & 4 & \\ \hline \end{array} \rightarrow \begin{array}{|c|c|c|c|} \hline & 2 & & \\ \hline 1 & 3 & 4 & 5 \\ \hline \end{array} \rightarrow \begin{array}{|c|c|c|c|} \hline & & 2 & \\ \hline 1 & 3 & 4 & 5 \\ \hline \end{array} \rightarrow \begin{array}{|c|c|c|c|} \hline & & & 2 \\ \hline 1 & 3 & 4 & 5 \\ \hline \end{array}$$

Proposition 15. For λ a partition of n , let $l = n - \lambda_1$ and fix k . Consider all μ with λ/μ a horizontal strip and $|\mu| = n - \lambda_1 + k$,

$$\sum_{\mu: |\mu|=l+k} d^\mu \leq \binom{l+k}{l-1} d_{\lambda/\lambda_1}$$

Reiner, Saliola and Welker [6] proved that there is a bijection between tableaux of λ and all desarrangement tableaux of the partitions μ for which λ/μ is a horizontal strip. The bijection can be described as: starting with $\hat{Q}$ a desarrangement tableaux of μ , with λ/μ a horizontal strip, append the cells of λ/μ as empty cells. Perform jeu-de-taquin slides into these empty cells in order from the leftmost to rightmost cell of λ/μ . As a consequence of λ/μ being a horizontal strip, the resulting tableau has the first $|\lambda/\mu|$ cells of the first row of λ empty. Filling the first row with $1, \dots, |\mu|$ and incrementing the rest by $|\lambda/\mu|$ gives a standard tableaux of λ .


Example 16. If $\lambda =$

, $\mu =$

 and $\hat{Q} =$

1	3	4
2	6	7
5		

. Then the jeu-de-taquin suggests



Reassigning the values, we get the following tableaux:

1	2	3	6
4	5	9	
7	8		


The following proposition describes the inverse process of the one introduced by Reiner, Saliola and Welker.

Proposition 17. To do the inverse of the above function, given a tableau, find the unique a, b so that the initial sequential entries in the first row are $1, 2, \dots, a$, the first column begins $1, a+1, \dots, a+b$, and the cell containing $a+b+1$ is not in the first column. If b is odd, remove $1, \dots, a-1$, perform jeu-de-taquin slides, and subtract $a-1$ from all entries. If b is even, remove $1, \dots, a$, perform jeu-de-taquin slides, and subtract a from all entries.

Example 18. Looking at

1	2	3	6
4	5	9	
7	8		

we see that $a = 3$ and $b = 1$. Therefore, we delete the entries 1 and 2 and we perform the jeu-de-taquin as follows:



Then we subtract $a = 2$ from the entries and we get exactly $\hat{Q} =$

1	3	4
2	6	7
5		

 which is what we started with in Example 16.

Proof of Proposition 17. For the inverse function, the resulting tableau after the deletion and jeu-de-taquin slides has first entry a if b is odd (or $a+1$ if b is even), increasing in the first column to $a+b$, and $a+b+1$ as the second entry of the first row. After removing $a-1$ if b is odd (or a if b is even) from all entries, this leaves the second entry of the first row as $b+2$ which is odd since b is odd (or $b+1$ which is even as b is even). This tableau is then a desarrangement tableau. $\square$

This construction on the inverse of the bijection allows for a good approximation of $\sum_{\mu: |\mu/\mu_\lambda|=k} d^\mu$. The below approximation echoes the bound on $\sum_{\lambda: \lambda_1=n-l} d_\lambda \leq \binom{n}{l} d_{\lambda/\lambda_1}$, and nearly recovers this when summing over k .

Proof of Proposition 15. We will bound the tableaux of λ that give rise to a partition of size $n - \lambda_1 + k$ after performing the inverse of the bijection in Proposition 17. Let $l = n - \lambda_1$. The number of blocks removed must be $n - (l + k) = n - l - k$. We will split this into cases based on the parity of b .

For b odd, this means the a in the inverse map is given by $a - 1 = \lambda_1 - k$. We need to count tableau with first row starting $1, \dots, a$ and first column starting $1, a + 1, \dots, a + b$. An upper bound for the number of such tableaux can be computed by selecting any $l - b$ elements of $a + b + 1, \dots, n$ as the entries that appear below the first row in any tableaux of $\lambda/[\lambda_1, 1^b]$, and the remaining entries in order in the remaining positions of the first row. This is at most

$$\binom{n - a - b}{l - b} d_{\lambda/\lambda_1} = \binom{l + k - b - 1}{l - b} d_{\lambda/\lambda_1} \leq \binom{l + k - b}{l - b} d_{\lambda/\lambda_1}$$

For b even, letting $a = \lambda_1 - k$, we need to count tableau with first row $1, \dots, a$ and first column starting $1, a + 1, \dots, a + b$. Using the same approximation technique, the number of such tableaux is at most:

$$\binom{n - a - b}{l - b} d_{\lambda/\lambda_1} = \binom{l + k - b}{l - b} d_{\lambda/\lambda_1}$$

Note that the final form of the bound is the same for both b odd and even, so summing over values of b of the binomial coefficients using the combinatorial identity $\sum_{i=1}^m \binom{n+i}{i} = \binom{n+m+1}{m}$ is:

$$\sum_{b=1}^l \binom{l + k - b}{l - b} \leq \binom{l + k}{l - 1}$$

Therefore,

$$\sum_{|\mu|=n-\lambda_1+k} d^\mu \leq \sum_{b=1}^l \binom{l + k - b}{l - b} d_{\lambda/\lambda_1} \leq \binom{l + k}{l - 1} d_{\lambda/\lambda_1}$$

which finishes the proof of the proposition. $\square$

3. $3/4n \log(n)$ UPPER BOUND

In this section, we present the proof of Theorem 1. To achieve the bound of $3/4n \log(n)$, its necessary to take advantage of how $\text{eig}(\lambda/\mu)$ decreases and d^μ increases as more blocks are added to μ from the minimal μ_λ . Although up to λ_1 blocks can be added to μ_λ , only the partitions μ with at most $\sqrt{(n - \lambda_1)n}$ added blocks will significantly contribute. This will reduce the contribution of $d_\lambda d^\mu$ to be considered from order $n^{2(n-\lambda_1)}$ to $n^{\frac{3}{2}(n-\lambda_1)}$ leading to a upper bound of $t = \frac{3}{4}n \log(n) + cn$.

To illustrate this, we will do as a special case the terms for $\lambda = [n - 1, 1]$, $\mu = [k, 1]$. The largest eigenvalue corresponds to $k = 1$ with multiplicity $n - 1$. If the mixing time were given only by the time it takes for $d_{[n-1,1]} d^{[1,1]} (\text{eig}([n - 1, 1]/[1, 1]))^{2t} = (n - 1) \left(1 - \frac{n+2}{n^2}\right)^{2t}$ to be exponentially small, the mixing time would be $\frac{1}{2}n \log(n)$. The spectrum of this walk is closely bunched around this eigenvalue, and so still contributes to the mixing time. If it were entirely bunched around this eigenvalue, the mixing time would be $n \log(n)$. That it is also spread out gives rise to the odd coefficient of $\frac{3}{4}$.

Proposition 19. For $t = \frac{3}{4}n \log(n) + cn$:

$$\sum_{k=1}^{n-1} d_{[n-1,1]} d^{[k,1]} (\text{eig}([n-1,1]/[k,1]))^{2t} = \sum_{k=1}^{n-1} (n-1) \left(1 - \frac{n+k^2+k}{n^2}\right)^{2t} \leq e^{-2c} + e^{-4c}$$

where $c > 0$

Proof. The tableaux of shape $[n-1, 1]$ all have $(1, 1)$ entry 1 and have $n-1$ further increasing entries in the first row, and so are completely determined by which of $2, \dots, n$ is in $(2, 1)$, so $d_{[n-1,1]} = n-1$. The number of desarrangement tableaux of $[k, 1]$ is 1 since the first column must be sequential up to an even number. The only option for the $(2, 1)$ entry is 2, so $d^{[k,1]} = 1$.

As (2) suggests, $\text{eig}([n-1, 1]/[k, 1]) = 1 - \frac{n+k^2+k}{n^2}$. To bound the contribution of these eigenvalues to the mixing time, we'll use the first $\sqrt{n}$ without using the $k^2 + k$ contribution. Once $k > \sqrt{n}$, the k^2 will be used.

$$\begin{aligned} \sum_{k=1}^{n-1} (n-1) \left(1 - \frac{n+k^2+k}{n^2}\right)^{2t} &\leq \sum_{k=1}^{n-1} (n-1) e^{-2t \frac{n+k^2+k}{n^2}} \\ &\leq \sum_{k=1}^{\sqrt{n}} (n-1) e^{-\frac{2t}{n}} + \sum_{k=\sqrt{n}+1}^{n-1} (n-1) e^{-\frac{4t}{n}} \\ &\leq n^{3/2} e^{-\frac{3}{2} \log n - 2c} + n^2 e^{-3 \log n - 4c} \leq e^{-2c} + e^{-4c} \end{aligned}$$

□

For all other λ , the d^μ are not uniform, and so understanding how they are distributed is essential in bounding $\sum_\mu d_\lambda d^\mu (\text{eig}(\lambda/\mu))^{2t}$. This is where we will use the bounds of sections 2.1 and 2.2. We are now ready to prove our main result:

Theorem 1. For the random to random card shuffling, if $t = \frac{3n}{4} \log n + cn$ with $c \geq 2$ and n is sufficiently large, then

$$\|P_{id}^{*t} - \pi\|_{T.V.} \leq e^{-c}.$$

Proof.

$$(4) \quad 4\|P^t - \pi\|_{TV}^2 \leq \sum_{l=1}^{n-1} \sum_{\lambda: \lambda_1=n-l} d_\lambda \sum_{k=0}^{\lambda_1} \left(\sum_{\mu: |\mu/\mu_\lambda|=k} d^\mu \right) (\text{eig}(\lambda/\mu))^{2t}$$

$$(5) \quad \leq \sum_{l=1}^{n-1} \sum_{\lambda: \lambda_1=n-l} \binom{n}{l} d_{\lambda/\lambda_1} \sum_{k=0}^{\lambda_1} \binom{k+l}{l-1} d_{\lambda/\lambda_1} (\text{eig}(\lambda/\mu))^{2t}$$

$$(6) \quad \leq \sum_{l=1}^{n-1} \sum_{\lambda: \lambda_1=n-l} \binom{n}{l} d_{\lambda/\lambda_1}^2 \sum_{k=0}^{n-l} \binom{k+l}{l-1} (\text{eig}(\lambda/\mu))^{2t}$$

$$(7) \quad \leq \sum_{l=1}^{n-1} \binom{n}{l} l! \left(\sum_{k=0}^l \binom{k+l}{l-1} \left(1 - \frac{l}{n}\right)^{2t} + \sum_{k=l+1}^{n-l} \binom{k+l}{l-1} \left(1 - \frac{ln+k^2+lk}{n^2}\right)^{2t} \right)$$

where in (5) we use Proposition 15 and in (6) we use that fact that $\sum_{\lambda: \lambda_1=n-l} d_{\lambda/\lambda_1}^2 = l!$. Here the bound will be split into cases (A) where $k \leq l$ where the improved bound of Proposition 11 on the eigenvalues does not apply, (B) and $k > l$.

For the terms of (A), note that $\sum_{k=0}^l \binom{k+l}{l-1} = \binom{2l+1}{l} \leq 2^l$ and $\binom{n}{l} l! \leq n^l$. The bound concludes by using $(1-x) \leq e^{-x}$, $n \geq 4$, and $2t = \frac{3}{2}n \log(n) + cn$.

$$(8) \quad (A) \leq \sum_{l=1}^{n-1} \binom{n}{l} l! \left(\sum_{k=0}^l \binom{k+l}{l-1} \left(1 - \frac{l}{n}\right)^{2t} \right)$$

$$(9) \quad \leq \sum_{l=1}^{n-1} n^l 2^l \left(1 - \frac{l}{n}\right)^{2t}$$

$$(10) \quad \leq \sum_{l=1}^{n-1} n^l 2^l e^{-\frac{3}{2}l \log(n) - 2lc}$$

$$(11) \quad \leq \sum_{l=1}^{n-1} e^{-\frac{1}{2}l \log(n/4) - l2c}$$

$$(12) \quad \leq \sum_{l=1}^{n-1} e^{-2lc}$$

$$(13) \quad \leq e^{-2c} \frac{e^4}{e^4 - 1}.$$

since $c \geq 2$.

Now we are ready to apply the same idea in order to bound (B). We will use Proposition 19 to handle the $l = 1$ case. We are going to further split the sum into two sums. For the first one will assume that $k \leq \sqrt{ln}$ and therefore we can neglect the $k^2 + k$ term. For the second one will have $k^2 \geq nl$. Let $t = \frac{3}{4}n \log n + cn$, then

$$(14) \quad (B) \leq \sum_{l=2}^{n-1} \binom{n}{l} l! \sum_{k=l+1}^{n-l} \binom{k+l}{l-1} \left(1 - \frac{l}{n} - \frac{k^2 + kl}{n^2}\right)^{2t}$$

$$(15) \quad \leq \sum_{l=2}^{n-1} \sum_{l+1 \leq k \leq \sqrt{ln}} n^l \frac{(k+l)^{l-1}}{(l-1)!} \left(1 - \frac{l}{n}\right)^{2t} + \sum_{l=1}^{n-1} \sum_{k > \sqrt{ln}} n^l \frac{(k+l)^{l-1}}{(l-1)!} \left(1 - \frac{2l}{n}\right)^{2t}$$

$$(16) \quad \leq \sum_{l=2}^{n-1} (\sqrt{ln} - l) n^l \frac{(\sqrt{nl} + l)^{l-1}}{(l-1)!} e^{-\frac{3}{2}l \log n - 2cl} + \sum_{l=1}^{n-1} n^{l+1} \frac{(2n)^{l-1}}{(l-1)!} e^{-3l \log n - 4cl}$$

To bound the first sum we will use Stirling's approximation

$$\left(\frac{l-1}{e}\right)^{l-1} \leq \frac{(l-1)!}{\sqrt{l-1}}$$

as well to simply,

$$(\sqrt{ln} - l) \frac{(\sqrt{nl} + l)^{l-1}}{(l-1)^{l-\frac{1}{2}}} e^{l-1} \leq n^{l/2} \left(\frac{l}{l-1}\right)^{l-\frac{1}{2}} \left(1 - \frac{1}{\sqrt{n}}\right) \left(\frac{1}{\sqrt{l}} + \frac{1}{\sqrt{n}}\right)^{l-1} e^{l-1} \leq 2^{3/2} n^{l/2} e^{l-1}$$

where the last inequality follows from two further bounds, since $l \geq 2$, $\left(\frac{l}{l-1}\right)^{l-\frac{1}{2}} \leq 2^{3/3}$ and since we have $l \geq 2$, $\frac{1}{\sqrt{l}} + \frac{1}{\sqrt{n}} \leq 1$.

$$\begin{aligned}
(16) &\leq \sum_{l=2}^{n-1} 2^{3/2} e^{l-1} e^{-2cl} + \sum_{l=1}^{n-1} \frac{1}{(l-1)!} e^{-4cl} \\
&\leq \frac{2^{3/2}}{e} e^{-2c} \frac{e^2}{e^2 - 1} + e^{-4c}
\end{aligned}$$

since $c \geq 2$.

Totaling our constants from the $l = 1$ case, (A), and (B) achieves a bound of

$$4\|P^t - \pi\|_{TV}^2 \leq e^{-2c} \left(1 + e^{-4} + \frac{1}{1 - e^{-4}} + \frac{2^{3/2}}{e} \frac{1}{1 - e^{-2}} + e^{-4} \right) \leq 4e^{-2c},$$

which finishes our proof. $\square$

4. RANDOM TO RANDOM WITH REPEATED CARDS

In this section, we briefly examine the case where the n cards of the deck are not necessarily distinct. Let $\nu = [\nu_1, \dots, \nu_m]$ be a partition of n . Then we can perform the random to random to a deck which has ν_i cards of type i , where $i = 1, \dots, m$. We then say that the deck has evaluation ν .

Theorem 2. *For the random to random on a deck with evaluation $\nu = [\nu_1, \dots, \nu_m]$ with $\nu_1 \geq \frac{n}{2}$, we have that*

(1) *For $t = \frac{3}{4}n \log n + cn$ with $c \geq 2$ and n sufficiently large, then*

$$\|P_{id}^{*t} - \pi\|_{TV} \leq \frac{1}{2} \left\| \frac{P_{id}^{*t}}{\pi} - 1 \right\|_2 \leq e^{-c}.$$

where $c > 2$.

(2) $t = \frac{n}{2} \log m + \frac{n}{4} \log n - cn$, then

$$\left\| \frac{P_{id}^{*t}}{\pi} - 1 \right\|_2 \geq \frac{1}{2} e^c.$$

where $c > 0$.

The proof is similar to the proof of Theorem 1, therefore we will quickly review what we need for this case. Diecker and Saliola [3] proved that the eigenvalues of this walk are given by

$$(17) \quad \text{eig}(\lambda/\mu) = \frac{1}{n^2} \left(\binom{n+1}{2} - \binom{|\mu|+1}{2} + \text{diag}(\lambda) - \text{diag}(\mu) \right)$$

with the restriction that $\lambda \succeq \nu$ which means that

$$\lambda_1 + \dots + \lambda_i \geq \nu_1 + \dots + \nu_i$$

for every $i = 1, \dots, m$. The multiplicity of such an eigenvalue is

$$\sum_{\substack{\lambda/\mu \text{ is a horizontal strip} \\ \lambda \succeq \nu}} K_{\lambda, \nu} d^\mu$$

where $K_{\lambda, \nu}$ is the number of semistandard Young tableaux (SSYT) of shape λ and evaluation μ and d^μ is the number of desarrangement tableaux of μ , just like before.

Proof of Theorem 2. The first part of Theorem 2 is easy to prove because $K_{\lambda,\nu} \leq d_\lambda$. Therefore, imitating the proof of Theorem 1 we get that for $t = \frac{3}{4}n \log n + cn$, then

$$\frac{1}{2} \left\| \frac{P_{id}^{*t}}{\pi} - 1 \right\|_2 \leq e^{-c}.$$

For the second part we only need to check the case where $\lambda = [n-1, 1]$. In this case, $K_{\lambda,\nu} = m$ and $d^\mu = 1$. Just like in Proposition 19, if $t = \frac{n}{4} \log m + \frac{n}{8} \log n - cn$, then

$$\sum_{k=1}^{n-1} m \left(1 - \frac{n+k^2+k}{n^2} \right)^{2t} \geq \sum_{k \leq \sqrt{n}-1} m \left(1 - \frac{2}{n} \right)^{2t} = (\sqrt{n}-1) m \left(1 - \frac{2}{n} \right)^{2t} \geq \frac{1}{2} e^{2c}$$

using that for $x \leq \frac{1}{2}$, $1-x \geq e^{-x(1+x)}$ and $c > 0$. $\square$

Acknowledgements. We would like to thank Franco Saliola for several enlightening conversations during the course of the project.

REFERENCES

- [1] Persi Diaconis and Laurent Saloff-Coste. Comparison techniques for random walk on finite groups. *Ann. Probab.*, 21(4):2131–2156, 1993.
- [2] Persi Diaconis and Mehrdad Shahshahani. Generating a random permutation with random transpositions. *Z. Wahrsch. Verw. Gebiete*, 57(2):159–179, 1981.
- [3] A. B. Dieker and F. Saliola. Spectral analysis of random-to-random markov chains. *arXiv:1509.08580*.
- [4] David A. Levin, Yuval Peres, and Elizabeth L. Wilmer. *Markov chains and mixing times*. American Mathematical Society, Providence, RI, 2009. With a chapter by James G. Propp and David B. Wilson.
- [5] Ben Morris and Chuan Qin. Improved bounds for the mixing time of the random-to-random insertion shuffle. *arXiv:1412.0070*, 2014.
- [6] Victor Reiner, Franco Saliola, and Volkmar Welker. Spectra of symmetrized shuffling operators. *Mem. Amer. Math. Soc.*, 228(1072):vi+109, 2014.
- [7] L. Saloff-Coste and J. Zúñiga. Refined estimates for some basic random walks on the symmetric and alternating groups. *ALEA Lat. Am. J. Probab. Math. Stat.*, 4:359–392, 2008.
- [8] Eliran Subag. A lower bound for the mixing time of the random-to-random insertions shuffle. *Electron. J. Probab.*, 18:no. 20, 20, 2013.
- [9] J.-C. Uyemura-Reyes. Random walks, semi-direct products and card shuffling. *ProQuest LLC, Ann Arbor, MI, 2002, Thesis (Ph.D.)-Stanford University. MR-2703300*.